



## King's Research Portal

DOI:

[10.1186/s12864-018-4997-y](https://doi.org/10.1186/s12864-018-4997-y)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Couto Alves, A., Glastonbury, C. A., El-Sayed Moustafa, J. S., & Small, K. S. (2018). Fasting and time of day independently modulate circadian rhythm relevant gene expression in adipose and skin tissue. *BMC GENOMICS*, 19, [659]. <https://doi.org/10.1186/s12864-018-4997-y>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

**Fasting and time of day independently modulate circadian rhythm relevant gene expression in adipose and skin tissue**

**Authors**

Alexessander Couto Alves<sup>1</sup>, Craig A. Glastonbury<sup>1</sup>, Julia S. El-Sayed Moustafa<sup>1</sup>, Kerrin S. Small<sup>1\*</sup>

<sup>1</sup>Department of Twin Research and Genetic Epidemiology, King's College London, St Thomas' Campus, Westminster Bridge Road, SE1 7EH London, United Kingdom.

\*Corresponding author: [kerrin.small@kcl.ac.uk](mailto:kerrin.small@kcl.ac.uk)

**Email addresses**

[Alexessander.Alves@gmail.com](mailto:Alexessander.Alves@gmail.com)

[craig@well.ox.ac.uk](mailto:craig@well.ox.ac.uk)

[julia.el-sayed\\_moustafa@kcl.ac.uk](mailto:julia.el-sayed_moustafa@kcl.ac.uk)

[kerrin.small@kcl.ac.uk](mailto:kerrin.small@kcl.ac.uk)

**Keywords**

Gene expression, fasting, circadian rhythm, eQTL, adipose, skin

## ABSTRACT

**Background:** Intermittent fasting and time-restricted diets are associated with lower risk biomarkers for cardio-metabolic disease. The shared mechanisms underpinning the similar physiological response to these events is not established, but circadian rhythm could be involved. Here we investigated the transcriptional response to fasting in a large cross-sectional study of adipose and skin tissue from healthy volunteers (N=625) controlling for confounders of circadian rhythm: time of day and season.

**Results:** We identified 367 genes in adipose and 79 in skin whose expression levels were associated ( $FDR < 5\%$ ) with hours of fasting conditionally independent of time of day and season, with 19 genes common to both tissues. Among these genes, we replicated 38 in human, 157 in non-human studies, and 178 are novel associations. Fasting-responsive genes were enriched for regulation of and response to circadian rhythm. We identified 99 genes in adipose and 54 genes in skin whose expression was associated to time of day; these genes were also enriched for circadian rhythm processes. In genes associated to both exposures the effect of time of day was stronger and in an opposite direction to that of hours fasted. We also investigated the relationship between fasting and genetic regulation of gene expression, including GxE eQTL analysis to identify personal responses to fasting.

**Conclusion:** This study robustly implicates circadian rhythm genes in the response to hours fasting independently of time of day, seasonality, age and BMI. We identified tissue-shared and tissue-specific differences in the transcriptional response to fasting in a large sample of healthy volunteers.

## Keywords

Gene expression, fasting, circadian rhythm, gene x environment, eQTL

## BACKGROUND

Time restricted and Calorie restricted diets are orthogonal interventions that constraint when and what is eaten. Calorie restriction is among the most reproducible life-prolonging interventions across taxa[1]. Intermittent and periodic fasting are two modalities of diet restriction that improve many cardio-metabolic risk factors and may increase healthspan in humans[2]. Surprisingly, time restricted produces strikingly similar results to calorie-restricted diet [3], despite only restricting the period when meals are consumed and not calorie intake. The reasons behind this have just started to be elucidated, but may involve the circadian rhythm[4]. The circadian rhythm controls a variety of physiological processes, including metabolism, sleep-wake cycle, and feeding. Central to these rhythms is the circadian clock, a transcription-translation feedback loop that can sustain an approximately 24-hour cycle. The central clock is located in the suprachiasmatic nuclei but other tissues also have clocks, e.g. heart, liver, kidney and adipose tissue[5]. The phase of the central clock is synchronized by light, but other factors such as mealtimes also entrain the phase of central and peripheral circadian clocks [6]. Both the absence and periodicity of mealtimes induced by intermittent fasting and time restricted diets may improve synchronization among central and peripheral clocks, which may lead to increased amplitude of clock gene expression[2]. Increased amplitude of the circadian rhythm oscillations is hypothesised to improve the overall health and cardio-metabolic status and disruption of circadian rhythm, e.g. shift work can lead to cardio metabolic conditions [2]. However, it is not clear how exactly this contributes to health status [7] or how to reconcile this with the pathogenesis of cardio metabolic disease, particularly the role of diet in connection with risk factors such as triglycerides, cholesterol, glucose, and insulin. Downstream pathways in peripheral tissues are plausible mediators of the circadian rhythm effect on health status, but the genes and the biological processes involved remain elusive.

One strategy to identify the molecular mechanisms mediating the effects of fasting on health is to identify fasting induced changes in gene expression across the body. Previous studies

of the relationship of gene expression with fasting and calorie restriction investigated adipose, blood, skeletal muscle, liver, and mucosa gene expression by contrasting controls with fasting cases [8] [9,10] or calorie restricted individuals[11][12][13][14][15,16][10,17–19], or by contrasting different times of the day [20]. However, none of these studies looked simultaneously at the effects of hours fasting and time of day on gene expression. Hence, it is not yet clear whether the effect of fasting is conditionally independent of the effect of time of day on circadian rhythms. Likewise, it is not yet clear if fasting and time of day response is independent of cardio metabolic risk factors. Other technical and design issues are also noteworthy, previous studies did not adjusted for the effects of seasonality, body mass index (BMI) and age, had smaller sample sizes (ranging from 4 to 311), used microarray or panels of genes, all of which may confound or bias the exploration of genes across the dynamic range of expression levels. Most importantly, skin transcriptome-wide gene expression response to hours fasting and time of day is yet uncharacterized.

Here we conducted an RNA-Seq gene expression association study on hours fasting adjusting for time of day, seasonality, BMI and age in adipose tissue and skin from 625 female twins from the TwinsUK registry. Adipose tissue plays a systemic role in metabolism, in the reproductive axis and in immune system homeostasis [21]. Skin is the largest organ and has an important systemic role on metabolism and the immune system [22] partly due to vitamin D synthesis. Circulating levels of vitamin D are associated with diabetes, cardiovascular disease, cancer, and osteoporosis [23]. Our aims are: 1) to map the pathways that independently respond to fasting and time of the day, to understand whether any of these pathways are common to both exposures, converging downstream into a common pathway; 2) to identify biological processes shared by both skin and adipose tissue as this may hint at details of the systemic organization of the response to fasting and time of day; 3) to identify novel genes responding to fasting independently of time of day and genes robustly replicating over the literature in multiple tissues in humans, non-humans and across taxa (i.e. both humans and non-humans) as this may hint at functional and/or conserved

pathways; 4) to assess the characteristics of common genetic variants regulating the expression of fasting responsive genes, by looking at expression quantitative trait loci (eQTL) and gene by environment (GxE) eQTL analyses.

## RESULTS

We analysed 625 female twins cross-sectionally embedded in the TwinsUK registry that satisfied the inclusion criteria (the exclusion was mostly driven by having all variables collected at the same time point). **Table 1** shows the characteristics of the individuals in this study. All individuals analysed are females with a mean age of 58 years and standard deviation (S.D.) of 9 years. The mean BMI was 27 kg.m<sup>-2</sup> and S.D. 5 kg.m<sup>-2</sup>. This is in agreement with the rest of the UK population[24] for the age bracket. The hours fasted ranged from 1-23h and the time of day when visits to the clinical facilities for the collection of the biopsy ranged from 08.30-14.30h. The visits occurred relatively uniformly between Monday to Friday in ~226 days out of 365 calendar days for nearly 20 months. A total of 250 women attended clinic between 8:30 and noon (AM) and 444 attended between noon and 14:30 (PM). Individuals attending clinic AM were asked to fast overnight and PM individuals were asked to not eat after 9AM. Self-reported hours fasting in AM, PM and overall varied considerably (**Figure 1**) suggesting social desirability bias was small. Abdominal Punch biopsies of subcutaneous adipose tissue (fat) and skin were extracted from a single punch biopsy taken during the clinical visit. RNA extraction, quantification and normalization were all performed as previously reported[25]. Expression levels were inverse-normal rank transformed to improve robustness of the analysis. We report our results at gene level for the best meta-exon, hereafter called exons unless otherwise stated. P values smaller than 10<sup>-16</sup> were rounded-up to P<10<sup>-16</sup> avoid numerical precision errors.

### Transcriptional response to fasting is more extensive in adipose tissue

To identify genes responding to hours fasting conditionally independent of time of the day, we regressed gene expression levels on hours fasting adjusting for time of the day, age, BMI, seasonality and other technical covariates using a mixed effects linear model (**Figure 2**). All exons significantly associated ( $P < 10^{-5}$  and  $FDR < 0.05$ ) with hours fasting, time of day and/or season are reported in **Supplementary Table S1**, Additional File 1. The response to hours fasting was more extensive and stronger in adipose than skin. We identified 367 genes responding to hours of fasting in adipose and 79 in skin (**Table 2 and Supplementary Table S1, Additional File 1**). The magnitude of the fasting effect was larger in adipose than skin in the 29 genes significant in both tissues (**Table 3**,  $|\beta_{fat}| - |\beta_{skin}| = 0.017$ , mean magnitude difference t-test  $P < 10^{-16}$ ). The response to time of day affected substantially less genes than fasting in both adipose (99) and skin (54), despite both exposures being assessed simultaneously (**Table 2**). The overall mean magnitude of the response to time of day was similar across tissues ( $|B| = 0.10$ ,  $P = 0.7$  and  $|\beta_{fat}| - |\beta_{skin}| = 0.001$ ,  $P > 0.8$ ) (**Table 3**). This suggests there are differences in which tissues respond to what systemic exposures and also differences in the magnitude of that response.

To identify tissue shared and tissue specific genes responding to hours of fasting and time of day, we looked directly at the overlap of significant associations by assessing the intersection of genes significant in both tissues (**Table 2**). We found 29 genes associated with hours fasting overlap across tissues (Overlap significance  $P < 10^{-16}$ , odds ratio (OR) = 18.9) and 17 for time of day (Overlap significance  $P < 10^{-16}$ , OR = 60.3). This corresponded to 38% and 31% of the genes associated with fasting and time of day respectively (**Table 2**). The common genes in either tissue are related to circadian clock or involved in the regulation of the circadian rhythm (**Figure 3**). The considerable overlap between tissues, particularly for hours fasting, suggest there is a systemic component in addition to the tissue-specific responses.

**Magnitude of the response to time of day is larger than hours fasting despite less genes associated**

To identify the overlap of genes responding to both fasting and time of day, we intersected the number of genes associated with both exposures in each tissue. (**Table 2**). The significance of the overlap was inferred with a fisher test as above. We found 71 genes in adipose (Overlap significance  $P < 10^{-16}$ ,  $OR = 37.1$ ) and 19 in skin (Overlap significance  $P < 10^{-16}$ ,  $OR = 67.4$ ) that respond to both hours fasting and time of day. This corresponds to 72% of genes associated with time of day in adipose and 35% in skin (**Table 2**). The substantial overlap of genes that respond to hours fasting conditionally independent of time of day in adipose suggests that these exposures elicit similar downstream processes.

To assess and compare the magnitude of the response to hours of fasting and time of day, we compared the average magnitude of the effect size (expression standard deviations/hour) of the exons significantly associated with each exposure in adipose and skin (**Table 3**). We found that the magnitude of adipose and skin response to time of day was around twice that of hours fasting ( $|\beta_{HF}| \approx 0.05$  and  $|\beta_{TOD}| \approx 0.1$ , difference in mean t-test  $P < 2 \times 10^{-16}$ ). Time of day effect is in the opposite direction of fasting for all exons significantly associated with both exposures (**Supplementary Table S1**, Additional File 1). These results suggest that the effect of time of day in gene expression is strong in these tissues, and may counteract the effect of fasting around the time clinical examinations occurred.

**Genes associated with hours fasting replicate among human tissues and across taxa**

To replicate our results, and identify genes that are associated with hours fasting consistently across human tissues and taxa, we conducted a literature search for gene expression papers of diet restriction (including fasting modalities) and queried the GenDR database for genetic manipulations in model organisms that cancel out or disrupt the life-extending effects



of diet restriction (**Supplementary Table 2**, Additional File 1). We looked at 3 types of replication: in human studies only, non-human studies only, or in both human and non-human studies. The later form of replication we termed “across taxa”. We assessed the replicability of each study by looking at the percentage of their genes replicated in other studies and likewise, we assessed the replicability of each gene by looking at the percentage of studies that reported the same gene (see methods). Among studies sampled from the literature, the median percentage of genes replicable in at least another study was 28.5% in humans and 57% in nonhumans. In both cases, studies on fasting and calorie restrictions show similar pattern of gene overlapping (**Supplementary Table S2**). In our study (**Supplementary Table S3**), 194 genes in adipose and 45 in skin replicated at least in one other study (Corresponding to 53% in adipose and 57% of genes in skin). In adipose 33 genes replicated across taxa, 24 in humans only, 137 in non-humans only (**Supplementary Table S3**, Additional File 1.). In skin 8 genes replicated across taxa, 3 in humans only, 34 in non-humans only (**Supplementary Table S3**, Additional File 1.). Four of the 19 genes associated to fasting in both tissues robustly replicate across taxa e: *PER1*, *NFIL3*, *CAMKK1*, *SPHK2*. *PPARG*, a key gene in adipogenesis and a Type 2 Diabetes drug target was implicated by our study in adipose tissue response to fasting, replicating previous results in model organisms. Our study implicates in the response to fasting for the first time 173 genes in adipose and 34 in skin. Novel genes in adipose include ageing-related *FOXO3*, and in skin *FOXN3-AS1*, a highly regulated non-coding gene with unknown function. Our replicability assessment shows that many genes responding to fasting in our study were replicated in multiple human tissues and are conserved across taxa, despite the multiplicity of designs and the fact that both studies in calorie restriction and fasting were included.

### **Genes associated with hours fasting are enriched for circadian rhythm**

We conducted a gene ontology enrichment analysis of the significant genes associated with hours fasting in adipose and skin using TopGO (**Supplementary Table S4**, Additional File

1.), ToppFun (**Supplementary Table S5**, Additional File 1.), and Ingenuity ( **Figure 4**, **Supplementary Table S6**, Additional File 1). TopGO implements multiple correction algorithms for the hierarchical structure of the GO. The ToppFun database is not based on entrez and as such maps a larger number of genes than TopGO. Ingenuity database has proprietary pathways that are manually curated. We found that circadian rhythm and gene expression regulation of circadian rhythm were the top enriched ontologies in both adipose and skin for both time of the day and hours fasting ( $P < 2.3 \times 10^{-05}$ , Fisher test BH FDR < 0.05). These results were robust to sensitivity analysis of the expression association P-value cutoff (**Supplementary Table S7, S8**, Additional File 1). The subset of genes associated with hours fasting in both skin and adipose (**Figure 3**) yielded a stronger enrichment for circadian rhythm than each tissue separately (BH FDR < 0.05), suggesting a systemic response. The identified genes annotated with circadian rhythm gene ontology and ingenuity circadian clock pathway were assessed for replication in other studies. Eight genes of the circadian clock associated with fasting were replicated in other studies, 1 across taxa and the remaining 7 only in non-humans: *PER1*, *PER2*, *NR1D1*, *ARNTL*, *CRY2*, *BHLHE40*, *BHLHE41*, *NPAS2*. Among the genes responsive to the circadian rhythm, three replicated across taxa (*KLF10*, *CREM*, *NFIL3*) and three only in non-human studies (*DBP*, *KLF9*, *PPARG*). Novel circadian rhythm genes associated with fasting include *PER3*, *CIART*, *UTS* and *PTGER4*. The overall set of replicable genes were highly enriched for circadian rhythm GO term (BH FDR <  $1.5 \times 10^{-3}$  in either adipose or skin).

In addition to the strong enrichment for circadian pathways, a range of other processes were also enriched in the genes associated to the two exposures (Supplementary Table S4-S6, Additional File 1). Focusing on the TopGO results, genes associated to fasting in adipose were enriched for glucose homeostasis ( $P = 1.6 \times 10^{-08}$ ), hormone secretion ( $P = 1 \times 10^{-08}$ ), regulation of vascular endothelial growth factor receptor signaling pathway ( $P = 6.4 \times 10^{-07}$ ). Genes associated to time of day were enriched for glucocorticoid signaling and response to lipid in both tissues, and to fat cell differentiation in skin only and a range of hormone and

ketone processes in adipose only. Previous studies in skeletal muscle identified genes in the Hypoxia and Inflammatory Response pathways. In our study these GO Terms are not significantly enriched after multiple test correction. However, 5 genes in the hypoxia GO Terms (BNIP3L, FAM162A, CBFA2T3, MYC, DDIT4) and 4 in the inflammatory response GO Term (PTGER4, C1QTNF3, ETS1, CRHBP) were significantly associated with hours fasting in our study, interestingly the hypoxia genes were not associated with time of day (Supplementary Table S1).

#### **Circadian clock pathway response to fasting not mediated or confounded by cardio metabolic risk factors, AM/PM visit and gene sequence parameters.**

To assess whether the enrichment for circadian genes in fasting and time of day were mediated or confounded by cardio metabolic risk factors, AM/PM interaction, residual confounding, gene sequence characteristics, and quantification method, we re-ran several models adjusting for several potential confounders (**Supplementary Table S9**, Additional File 1) and repeated the enrichment analysis using TopGO in adipose and skin (**Supplementary Table S10**, Additional File 1). We found that circadian rhythm related pathways were always the top enriched GO term regardless of the model tested (**Supplementary Table S10**, Additional File 1). Adjusting for glucose and total cholesterol (which together with BMI and age explains hours fasting association with triglycerides, insulin, TC, LDL-C and HDL-C levels, see methods and **Supplementary Table S11**, Additional File 1) did not change the ranking of the top gene ontologies, which means enrichment is not mediated or confounded by these metabolic pathways. We stratified the data into AM and PM samples and the enrichment was still present in PM (AM sample was smaller). As restricting the sample to individuals that attended PM visits controls the effect of the time of the day, residual confounding with this covariate can be excluded as a potential explanation for the association of hours fasting with gene expression levels. We then adjusted Gene P-values for the number of exons, gene length, GC content, etc (see methods) but this did not change the result. We concluded from this analysis that circadian

rhythms in gene expression are influenced by hours fasting independently of cardio metabolic risk factors and not confounded with the technical factors and parameters that we can sensibly implicate in the expression response of these tissues.

## **Relationship between fasting and genetic regulation of gene expression**

To investigate whether change in expression of the genes responding to fasting and time of day may have a role in disease and other phenotypes, we investigated whether *cis*-eQTLs of the exposure-responsive genes are coincident with complex trait GWAS signals. As GWAS-coincident *cis*-eQTLs place modulation of expression of these genes on the molecular pathway between the genetic variant and trait, it is plausible that fasting or time of day-induced changes in expression could also impact the linked trait. Previous *cis*-eQTL scans of this dataset[25] identified 195 independent regulatory variants at 195 of the 553 exposure-responsive genes; we identified complex trait GWAS signals coincident with 11 of these 165 regulatory variants (or proxies  $R^2 > 0.8$ ) with the Phenoscanner portal [26] (**Table 4**, **Supplementary Table S12**, Additional File 1). Five genes responsive to hours of fasting (*CTPS1*, *DNM3*, *HEMK1*, *SLA3R2*, *TRIM27*, *TSEN15*) are regulated by *cis*-eQTLs coincident with height GWAS signals (**Table 4**). Fasting responsive genes were also regulated by variants associated to Crohn's disease (*PTGER4*), Rheumatoid arthritis (*PHLDB1*), Lung Function (*ADAM19*) and with cardiac troponin (*TNNT2*). *NFAIP8*, which is responsive to time of day in adipose tissue, is regulated by a Waist-hip ratio associated variant.

The response to external environments is not always uniform across individuals, and in some cases can be modified by the genotype of an individual. To identify common regulatory variants that modulate the response to hours of fasting or time of day we conducted a Gene x Hours-of-Fasting and Gene x Time-of-day *cis*-eQTL scan in each tissue. Overall, we identified 83 GxE interactions at an FWER < 0.05, with the largest number identified for Gene x Hours-of-Fasting in adipose (**Supplementary Table S13**, Additional File 1). Looking

at the 553 genes whose expression levels were associated to one of the exposures, we identified significant interactions at 15 genes (**Table 5**); 12 with hours fasting in adipose, one with time of day in adipose and one with time of day in skin (Three examples are plotted in **Figure 5**). The significant overlap with exposure-responsive genes ( $OR=7.6$ ,  $P=1.3 \times 10^{-08}$ ) suggest that genetic background may define the individual response to fasting and time of day at certain genes.

## DISCUSSION

We conducted an association study between gene expression and hours fasting, adjusting for time of day, seasonality, BMI, age and other technical covariates in a large sample of human adipose and skin biopsies. We identified 417 genes associated with hours of fasting in adipose and skin, which partly overlap with 137 genes associated to time of day in these samples. More than 50% of genes responding to fasting in our study were corroborated by a literature analysis of other gene expression studies on fasting and calorie restriction in other species and other human tissues. The response to hours fasting was more extensive in number of genes and stronger in magnitude in adipose tissue than skin, which is not unexpected given the central role of adipose tissue in metabolism. Interestingly, in genes associated to both exposures, the effect of time of day was stronger and in an opposite direction to the effect of hours of fasting, and thus may counteract the effect of fasting in some genes during the observed time window. This effect is consistent with clinical trials of the physiological response to meal times[27] suggesting improved cardio metabolic risk factors and weight management for those consuming most calories in the first half of the day. This finding may have consequences for deciding the optimal period of the day for fasting and planning of biomedical interventions.

Fasting responsive genes were enriched for circadian rhythm and circadian clock genes among other biological processes. The enrichment for circadian rhythm did not subside after

sensitivity analysis, re-running the associations conditioning on cardio-metabolic risk factors, or when restricting to samples collected in the afternoon. Genes with transcriptional responses to both exposures in both tissues were even more highly enriched for circadian rhythms and circadian clock suggesting this is systemic component of the fasting response in adipose and skin of humans. A role of gene expression circadian rhythm in time-restricted diets has been hypothesized based on animal models and *in vitro* studies of human materials[2,3] but the case for hours fasting is not yet fully established, particularly it was not clear whether the association is not confounded with time of the day, seasonality, and risk factors involved in the pathogenesis and pathophysiology of cardio metabolic disease. Our results contribute to this discussion by providing data from an observational study in multiple human tissues with approximately 625 individuals, the largest so far. Our results consolidate the theory that response to fasting and time of day share common pathways[4], particularly the circadian clock. The similarity of the beneficial effects on health of both time restricted diet and fasting might be underpinned by this common pathway, independently of known risk factors of the pathogenesis of cardio metabolic disease, such as cholesterol and triglycerides.

Regulation of genes response to time of day and hours fasting was tissue specific, with greater proportion of genes responding in fat, and interestingly the percentage overlap of genes responding to both exposures also larger in fat. We also found a larger proportion of regulatory variants interacting with hours fasting in fat; showing that expression plasticity in response to fasting is tissue specific. This support the notion that gene expression regulation is tissue-specific[28] and that tissue-specific regulatory variants may influence the onset of conditions affecting a particular tissue type. An idea supported by the fact that several exposure-responsive genes had regulatory variants overlapping GWAS signals of complex diseases.

Some limitations are important to mention, this is an observational twin-study of females with average age of 58 years and the hours fasting was self-reported which may be subject to error, recall bias, social desirability bias and self-selection bias. We removed individuals that fasted more than 23h to avoid recall bias. Self-selection bias in hours of fasting, e.g. for medical reasons, is small as this is a relatively healthy sample. As the time of the visit was allocated more or less haphazardly, this exposure will be less affected by confounding. The study may have missed circadian rhythm genes with acrophase centred outside the collection interval or that respond less vigorously to time of day in these tissues. The circadian rhythm genes (particularly those in the canonical circadian clock) were the most significantly associated genes with time of the day (and with fasting duration), which indicate the response of at least these circadian rhythm genes was vigorous and centred in the sampling interval. In order to identify further circadian rhythm responsive genes, the biopsy collection needs to be extended over a broader time interval, including the night hours. As this is an observational study, individuals that fasted long hours tend to fast overnight. Consequently, there is a chance of a putative interaction or confounding between duration and modality of fasting, i.e. between fasting overnight and fasting long hours. Nevertheless, the effects can still be ascribed to fasting regardless of the underlying characteristic mediating the effect. While our sample size ( $N \sim 625$ ) is well powered to detect associations between gene expression and our exposures, it is relatively small for detection of genetic interaction effects, which limits the power of the GxE scan. GxE interactions are difficult to detect in humans, but recently have been detected on regulation of gene expression in several studies[29][30,31] where the reduced search space, large effect size and in some cases use of cell lines allows greater power to identify interactions. As such, we report the results of the GxE scan, but note these results will need to be tested in other datasets to determine their robustness. Some strengths of the study are also worth mention. As we carefully selected only external exposures and avoided endogenous or intermediate phenotypes, the study is more robust to collider bias and confounding.

## CONCLUSIONS

This study identified 417 genes whose expression is associated to hours fasting independently of time of day and seasonality, two important confounders of circadian rhythms. 178 of these genes have not previously been implicated in the response to fasting, and another 171 have not been linked to fasting in humans before. We identified tissue-shared and tissue-specific differences in the transcriptional response to fasting, and identified a core set of circadian rhythm genes that responding to both fasting and time of day in both adipose and skin tissue. This study contributes to the characterization of the transcriptional response to fasting in humans. Gene expression studies like this provide unique opportunities to unveil the molecular mechanisms of the response to exposures and therefore contributes to the identification of targets for therapy or drug development, and the conditions under which such targets vary.



## **METHODS**

### **Selection of individuals and embedded design**

Twin pairs from the Twins UK registry were invited to clinical visit by telephone call and allocated to available time slots in the weekdays. The allocations happened haphazardly but subject to both twin pairs availability. Twins are routinely followed up over the years and multiple observations for some variables are available. We embedded a cross-sectional study on the subset of individuals whose materials for RNA extraction, hours fasting and clinical markers were collected on the same day that the anthropometric measurements were taken, with the exception of DNA which was collected previously. The rationale for using cross-sectional design follows directly from the time-dependent nature of the gene expression response to fasting and time of the day. We collected data over a period of 6 hours which is 50% of the elapsed time between maximum and minimum amplitude (average 12 hours for a period of 24 hours). For the genes with acrophase at 8h30m or 14h30m the study can detect up to 50% of the total peak-to-peak difference (the maximum signal). We further restricted our analysis to individuals that fasted less than 23 hours to avoid recall bias. Out of the ~850 individuals in all tissues, 625 individuals that meet all above criteria remained. Twins visits to the clinic were made over a period of ~600 days. The analyses of the collected materials were also performed at different times and in different batches. We recorded these dates and batches for investigation of potential confounders.

### **Tissue Collection**

The study sample included 856 female twins who are a part of the TwinsUK registry and are all of European ancestry. Punch biopsies of subcutaneous adipose tissue from a sunlight-protected area of the stomach adjacent and inferior to the umbilicus were obtained from consenting individuals. Skin from the punch biopsy was then dissected to separate it from adipose tissue, and both samples were weighed and immediately frozen with liquid nitrogen. All the procedures followed were in accordance with the ethical standards of the St. Thomas

Research Ethics Committee (reference 07/H0802/84) at St. Thomas Hospital in London. Volunteers gave informed consent and signed a consent form before the biopsy procedure. Volunteers were supplied with an appropriate detailed information sheet regarding the research project and biopsy procedure by post before attending the biopsy.

## **Genotyping and Imputation**

Genome-wide SNP data for the TwinsUK individuals were generated as previously described. [32] In short, TwinsUK samples were genotyped on a combination of platforms (HumanHap300, HumanHap610Q, and 1M-Duo Illumina arrays). Quality control and merging of the array datasets has previously been described in detail [32]. The cleaned data were pre-phased with IMPUTE2 with no reference panel and were then imputed into the 1000 Genomes phase 1 reference panel (interim, data freeze accessed November 10, 2010; the 1000 Genomes Project Consortium 2012). Variants with an INFO score >0.8 on all platforms and a MAF >5% were retained for analysis.

## **RNA-Seq, Exon Quantification and Normalization**

RNA-seq data were generated, quantified and normalized as previously described [25]. In brief, samples were prepared for sequencing with the unstranded Illumina TruSeq sample preparation kit and sequenced on a HiSeq 2000 machine. The 49 bp paired-end reads were aligned to the UCSC Genome Browser GRCh37 reference genome with the Burrows-Wheeler Aligner. GENCODE v.10 was used to annotate genes. Samples were excluded if they failed to have more than 10 million reads map to known exons or if the sequence data did not correspond to actual genotype data. To quantify exons, all overlapping exons of a gene were merged into one meta-exon. We counted reads as mapping to a given meta-exon if either of its start or end coordinates overlapped a meta-exon boundary. All read-count quantifications were corrected for variation in sequencing depth between samples by normalizing the number of reads to the median number of well-mapped reads. We only used

exons that were quantified in more than 90% of the individuals. Exon expression values were rank-based inverse normal transformed for downstream analysis.

#### **Covariates and Clinical Markers recorded**

We considered the following previously collected anthropometric and metabolic markers: total cholesterol, total triglycerides, HDL cholesterol, glucose, and insulin. We also collect information on potential technical confounders, including the day of the year (1-365, for seasonality), age (years), day of the study (0-600 days), day the material was analysed in the lab (factor), day RNA-Seq was conducted (factor). The following RNA-Seq technical covariates were collected for each individual: GC content, median insert size, primer index.

#### **Transcriptome-wide association analysis of expression levels**

Transcriptome-wide analysis were conducted using mixed effects models implemented in lmer package of R. The model used for all analysis unless otherwise stated is:

$$y_i = XB + Zu + \epsilon_i$$

Where  $y_i$  is a vector of  $n$  individuals with the expression levels of exon  $i$ ,  $X$  and  $Z$  are the design matrices of the fixed and random effects containing  $n$  individuals in the rows. The fixed effects include: hours fasting, time since last meal up to start of the clinical visit expressed in hours; *BMI*, the body mass index; age, years since born; time of day, local time of the clinical visit expressed in hours since midnight; Study day, number of days since the study started up to the day of the visit; the day of the year (ranging from 1-365); percentage of GC content in the aligned reads of the RNA-Seq data; the median insert size. The random effects were: Primer index; Date RNA-Seq experiment was conducted for this sample; Family identifier; Clonality identifier defined uniquely for each DNA sequence, such that a pair of MZ twins have the same id (because they have the same DNA) but each twin from a DZ pair have unique ids. These technical factors are an intrinsic part of the study design and

should be included in the model. In addition, gene expression in this sample and others has been shown to be associated with age (Vinuela et al, 2018), BMI (Glastonbury et al, 2016) and season (Dopico et al, 2015). As dawn time is a major cue for the synchronization of the circadian rhythm, we adjust for seasonality to avoid confounding with hours fasting and time of the day.

#### **Statistical significance of the gene expression association study**

As expression levels are highly dependent, we corrected for multiple testing by applying the Benjamini Hochberg algorithm to hours fasting and time of the day (sample biopsy) in skin and fat and selecting the most stringent cut-off from all analyses ( $P < 1 \times 10^{-5}$ ) that guarantees a Benjamini Hochberg  $FDR \leq 0.05$  in all tissues and all exposures assessed.

#### **Replication of genes in the literature**

We conducted two systematic searches in pubmed for papers reporting gene expression analysis of the response to fasting and calorie restriction separately in human and nonhuman mammals. As the automatic annotation of papers with MESH tags is not perfect, we enriched the search with the following strategies: i) retrieving all papers with “gene expression” keyword, selecting all relevant studies among the most recent top 100 papers; ii) using the same search string on google scholar, mendeley, and google. Once all relevant papers were identified, we extracted the significant genes reported in these studies, we converted probes, gene ids or gene symbols to ensembl ids using ensembl biomaRt. Studies reporting EST we used Genbank batch query followed by scripts developed in house to

506 extract the RefSeq Id (e.g. NM\_91388) and then converted these using the same process  
507 described above. Swissprot ids were converted to ensembl id using Swissprot tools. Non-  
508 human genes were mapped to ensembl ids using the strategies above and then mapped to  
509 human orthologues using ensembl bioma. Once all genes for all species were converted to  
510 human orthologues, we calculated the gene overlap between each study, including our  
511 study. We analysed two metrics to assess study replicability: Proportion of overlap between  
512 gene lists of study A and B was defined as  $|A \cap B| / \min(|A|, |B|)$ ; The percentage of the study  
513 genes replicated was defined as the number of genes that overlaps with at least another  
514 study divided by the total number of genes of study  $A_i$ :  $| \cup_j (A_i \cap A_j) | / |A_i|$  for all  $j \neq i$

515

## 516 **Gene ontologies enrichment**

517 We conducted gene ontology term enrichment using definitions of biological processes GO  
518 terms downloaded from R package org.Hs.eg.db, corresponding to a total number of 5828 in  
519 fat and 5992 in skin GO terms. All analyses restricted to terms <500 genes. GO term  
520 enrichment was independently conducted on genes significantly associated with Hours  
521 fasting and Time of the day using the TopGO package from R. We implemented all  
522 enrichment tests present in the package. GO terms that pass bonferroni correction ( $P < 10^{-5}$ )  
523 either on Fisher Elimination or Fisher methods were considered significant but results for all  
524 algorithms are presented (mostly consistent across all methods). We repeated the  
525 enrichment analysis using Ingenuity and ToppFun due to quality of the annotation database.  
526 Except for ToppFun, all analyses were conducted considering only the genes expressed in  
527 the tissue. The gene annotations with the GO circadian rhythm and ingenuity circadian clock  
528 pathway were further confirmed against the literature.

529

### 530 **Sensitivity of GO term enrichment to P-value cut-off**

531 We tested sensitivity of results to the significance cut-off by applying a more stringent cutoff  
532 on P-value and repeating the enrichment analysis. We considered two cutoffs: i)  $P < 2 \times 10^{-6}$ ,  
533 equivalent to at least Benjamini-Yekutieli  $FDR \leq 0.05$  in all tissues and all exposures and ii)  
534  $P < 5 \times 10^{-7}$  corresponding to Bonferroni correction of at least 100,000 tests, which is  
535 approximately the total number of exons in fat and skin (114-116,000).

536

### 537 **Sensitivity to confounders of the power to detect expression associations**

538 In order to test sensitivity to confounders of the power to detect associations with gene  
539 expression, we conducted a separate analysis correcting the P-value of the associations of  
540 exons with hours fasting by regressing the logit transformed P value on the percentage of  
541 GC content in the DNA sequence of the gene, the gene length in bp, the number of  
542 isoforms reported for this gene, the number of meta-exons, the number of independent  
543 meta-exons (calculated with Ji and Li method using eigendecomposition of expression levels  
544 of all meta-exon in a gene) and adjusted for the following random intercept effects: Gene  
545 type, gene status and gene annotation database as defined by GenCode version 10. The  
546 residuals of the regression model were then back-transformed and used to ascertain the  
547 total number of significant genes and the respective gene ontologies enriched.

548

### 549 **Identification of potential cardiometabolic risk factors mediating or confounding the** 550 **expression response to fasting**

551 We regressed Glucose, TC, TG, HDL-C on hours fasting adjusting for age, age<sup>2</sup>, BMI, time  
552 of day, and the following random effects: Family, Clonality, Biopsy Date using a mixed effects  
553 model. We identified the subset of variables that are potential mediators or confounders of  
554 the effects of hours of fasting in the circulating levels of these cardiometabolic markers by i)

identifying those markers associated with hours fasting, ii) discarding all markers whose association with hours fasting was abrogated after adjusting for a significant cardiometabolic marker.

### **Correction for population structure and eQTL analysis**

We correct population structure and twin relatedness using ProbABEL [33] two-step residual-outcome mixed models approach. First, we regress expression levels on technical covariates and kinship matrix using a mixed effects model.

$$\text{Expression} = XB + u + y, y \sim N(0, S), u \sim N(0, \sigma_a K)$$

The fixed effect  $X$  matrix have GC content and Insert size mode variables in the columns and individuals in the rows. The random effects  $u$  matrix have the primer index and date of RNAseq analysis as a proxy for batch effects in the columns. The kinship matrix  $K$  was estimated from imputed genotypes with INFO>0.80 and MAF>5% using GEMMA as previously described [31]. The residuals of the regression  $y$  is the outcome of this step. This regression is repeated for the expression of all genes or exons. The outcome of this step is a matrix of residuals with exons in the columns and individuals in the rows  $Y = \{y_1, y_2, \dots, y_m\}$ . In the second step we regress the expression residuals  $y$  on SNP and additional covariates using the linear regression implemented in MatrixEQTL. Expression principal components were estimated from the expression residuals  $Y$  matrix using prcomp implemented in R. We analysed two eQTL models corresponding to the hours fasting ( $T_f$ ) model (eq. 1) and local time of visit ( $T_v$ ) model (eq. 2). Notice that  $y$  are the expression residuals corrected for population structure using the imputed data as well as corrected for technical covariates using both fixed and random effects.

### **Statistical significance of GxE eQTL**

582 We conducted a permutation test by fixing all covariates and randomizing the expression  
583 levels. The permutation was synchronized across all exons to preserve transcriptome-wide  
584 correlation structure. On each permutation, we save the smallest p-value from all SNPs  
585 associated with each exon. In order to correct for the correlation structure between exons  
586 from the same gene, we grouped the exon permutation p-values by gene to produce a more  
587 stringent gene-level correction of p-values. The permuted p-values were compared to the  
588 association p-value to obtain the FWER.

589

#### 590 **Abbreviations**

591 G x E: Gene by environment interaction

592 eQTL: Expression quantitative trait loci

593 OR: Odds ratio

594 BMI: Body mass index

595 AM: Ante meridiem

596 PM: Post meridiem

597 TOD: Time of day

598



## **DECLARATIONS**

### **Ethics approval and consent to participate**

This project was approved by the ethics committee at St Thomas' Hospital London, where all the biopsies were carried out. Volunteers gave informed consent and signed an approved consent form prior to the biopsy procedure. Volunteers were supplied with an appropriate detailed information sheet regarding the research project and biopsy procedure by post prior to attending for the biopsy.

### **Consent for publication**

Not applicable

### **Availability of data and material**

TwinsUK RNAseq data is available from EGA (Accession number: EGAS00001000805; <https://www.ebi.ac.uk/ega/studies/EGAS00001000805>). TwinsUK genotypes are available upon application to the TwinsUK cohort – application procedures are described here: <http://www.twinsuk.ac.uk/data-access/submission-procedure-2/> .

### **Competing interests**

None declared

### **Funding**

This study was supported by MRC Project Grant (MR/L01999X/1) to K.S.S. TwinsUK is funded by the Wellcome Trust, Medical Research Council, European Union, the National Institute for Health Research (NIHR)-funded BioResource, Clinical Research Facility and Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust in

625 partnership with King's College London. . This project was enabled through access to the  
626 MRC eMedLab Medical Bioinformatics infrastructure, supported by the Medical Research  
627 Council [grant number MR/L016311/1]

628

#### 629 **Authors' contributions**

630 Study conception and design: ACA

631 Acquisition and pre-processing of data: ACA, KSS, CG, JSESM

632 Analysis and interpretation of the data: ACA, KSS

633 Drafting manuscript: ACA, KSS

634 Critical revision: KSS, CG, JSESM, ACA

635 All authors read and approved the final manuscript

636

#### 637 **Acknowledgements**

638 Authors wish to thank all twins that kindly participated in this study.

639

640

641

642

643

644

645

646

647

648

## References

1. Kenyon CJ. The genetics of ageing. *Nature*. 2010;464: 504–512.
2. Longo VD, Panda S. Fasting, Circadian Rhythms, and Time-Restricted Feeding in Healthy Lifespan. *Cell Metab*. 2016;23: 1048–1059.
3. Panda S. Circadian physiology of metabolism. *Science*. 2016;354: 1008–1015.
4. Froy O. Circadian Rhythms, Aging, and Life Span in Mammals. *Physiology* . 2011;26: 225–235.
5. Kiehn J-T, Tsang AH, Heyde I, Leinweber B, Kolbe I, Leliavski A, et al. Circadian Rhythms in Adipose Tissue Physiology. *Compr Physiol*. 2017;7: 383–427.
6. Mohawk JA, Green CB, Takahashi JS. Central and Peripheral Circadian Clocks in Mammals. *Annu Rev Neurosci*. 2012;35: 445–462.
7. Wei M, Brandhorst S, Shelehchi M, Mirzaei H, Cheng CW, Budniak J, et al. Fasting-mimicking diet and markers/risk factors for aging, diabetes, cancer, and cardiovascular disease. *Sci Transl Med*. 2017;9. doi:10.1126/scitranslmed.aai8700
8. Wijngaarden MA, van der Zon GC, van Dijk KW, Pijl H, Guigas B. Effects of prolonged fasting on AMPK signaling, gene expression, and mitochondrial respiratory chain content in skeletal muscle from lean and obese individuals. *Am J Physiol Endocrinol Metab*. 2013;304: E1012–21.
9. Wijngaarden MA, Bakker LEH, van der Zon GC, 't Hoen PAC, van Dijk KW, Jazet IM, et al. Regulation of skeletal muscle energy/nutrient-sensing pathways during metabolic adaptation to fasting in healthy humans. *Am J Physiol Endocrinol Metab*. 2014;307: E885–95.
10. Bouwens M, Afman LA, Müller M. Fasting induces changes in peripheral blood mononuclear cell gene expression profiles related to increases in fatty acid beta-oxidation: functional role of peroxisome proliferator activated receptor alpha in human peripheral blood mononuclear cells. *Am J Clin Nutr*. 2007;86: 1515–1523.
11. Dahlman I, Linder K, Arvidsson Nordström E, Andersson I, Lidén J, Verdich C, et al. Changes in adipose tissue gene expression with energy-restricted diets in obese women. *Am J Clin Nutr*. 2005;81: 1275–1285.
12. Armenise C, Lefebvre G, Carayol J, Bonnel S, Bolton J, Di Cara A, et al. Transcriptome profiling from adipose tissue during a low-calorie diet reveals predictors of weight and glycemic outcomes in obese, nondiabetic subjects. *Am J Clin Nutr*. 2017;106: 736–746.
13. Capel F, Viguerie N, Vega N, Dejean S, Arner P, Klimcakova E, et al. Contribution of energy restriction and macronutrient composition to changes in adipose tissue gene expression during dietary weight-loss programs in obese women. *J Clin Endocrinol Metab*. 2008;93: 4315–4322.
14. Lam YY, Ghosh S, Civitarese AE, Ravussin E. Six-month Calorie Restriction in Overweight Individuals Elicits Transcriptomic Response in Subcutaneous Adipose Tissue That is Distinct From Effects of Energy Deficit. *J Gerontol A Biol Sci Med Sci*. 2016;71: 1258–1265.

- 691 15. Campbell KL, Foster-Schubert KE, Makar KW, Kratz M, Hagman D, Schur EA, et al.  
692 Gene expression changes in adipose tissue with diet- and/or exercise-induced weight  
693 loss. *Cancer Prev Res* . 2013;6: 217–231.
- 694 16. Mercken EM, Crosby SD, Lamming DW, JeBailey L, Krzysik-Walker S, Villareal DT, et  
695 al. Calorie restriction in humans inhibits the PI3K/AKT pathway and induces a younger  
696 transcription profile. *Aging Cell*. 2013;12: 645–651.
- 697 17. Hietaniemi M, Jokela M, Rantala M, Ukkola O, Vuoristo JT, Ilves M, et al. The effect of a  
698 short-term hypocaloric diet on liver gene expression and metabolic risk factors in obese  
699 women. *Nutr Metab Cardiovasc Dis*. 2009;19: 177–183.
- 700 18. Pendyala S, Neff LM, Suárez-Fariñas M, Holt PR. Diet-induced weight loss reduces  
701 colorectal inflammation: implications for colorectal carcinogenesis. *Am J Clin Nutr*.  
702 2011;93: 234–242.
- 703 19. Gille D, Zangger N, Soneson C, Bütikofer U, Delorenzi M, Schwander F, et al. Caloric  
704 dose-responsive genes in blood cells differentiate the metabolic status of obese men. *J*  
705 *Nutr Biochem*. 2017;43: 156–165.
- 706 20. Loboda A, Kraft WK, Fine B, Joseph J, Nebozhyn M, Zhang C, et al. Diurnal variation of  
707 the human adipose transcriptome and the link to metabolic disease. *BMC Med*  
708 *Genomics*. 2009;2: 7.
- 709 21. Kershaw EE, Flier JS. Adipose tissue as an endocrine organ. *J Clin Endocrinol Metab*.  
710 2004;89: 2548–2556.
- 711 22. Di Meglio P, Perera GK, Nestle FO. The multitasking organ: recent insights into skin  
712 immune function. *Immunity*. 2011;35: 857–869.
- 713 23. Holick MF. Vitamin D: importance in the prevention of cancers, type 1 diabetes, heart  
714 disease, and osteoporosis. *Am J Clin Nutr*. 2004;79: 362–371.
- 715 24. Bhaskaran K, Douglas I, Forbes H, dos-Santos-Silva I, Leon DA, Smeeth L. Body-mass  
716 index and risk of 22 specific cancers: a population-based cohort study of 5·24 million UK  
717 adults. *Lancet*. 2014;384: 755–765.
- 718 25. Buil A, Brown AA, Lappalainen T, Viñuela A, Davies MN, Zheng H-F, et al. Gene-gene  
719 and gene-environment interactions detected by transcriptome sequence analysis in  
720 twins. *Nat Genet*. 2015;47: 88–91.
- 721 26. Staley JR, Blackshaw J, Kamat MA, Ellis S, Surendran P, Sun BB, et al. PhenoScanner:  
722 a database of human genotype-phenotype associations. *Bioinformatics*. 2016;32: 3207–  
723 3209.
- 724 27. Jakubowicz D, Barnea M, Wainstein J, Froy O. High caloric intake at breakfast vs.  
725 dinner differentially influences weight loss of overweight and obese women. *Obesity* .  
726 2013;21: 2504–2512.
- 727 28. López-Maury L, Marguerat S, Bähler J. Tuning gene expression to changing  
728 environments: from rapid responses to evolutionary adaptation. *Nat Rev Genet*. 2008;9:  
729 583–593.
- 730 29. Fairfax BP, Makino S, Radhakrishnan J, Plant K, Leslie S, Dilthey A, et al. Genetics of  
731 gene expression in primary immune cells identifies cell type-specific master regulators  
732 and roles of HLA alleles. *Nat Genet*. 2012;44: 502–510.

- 733 30. Moyerbrailean GA, Richards AL, Kurtz D, Kalita CA, Davis GO, Harvey CT, et al. High-  
734 throughput allele-specific expression across 250 environmental conditions. *Genome*  
735 *Res.* 2016;26: 1627–1638.
- 736 31. Glastonbury CA, Viñuela A, Buil A, Halldorsson GH, Thorleifsson G, Helgason H, et al.  
737 Adiposity-Dependent Regulatory Effects on Multi-tissue Transcriptomes. *Am J Hum*  
738 *Genet.* 2016;99: 567–579.
- 739 32. Suhre K, Shin S-Y, Petersen A-K, Mohny RP, Meredith D, Wägele B, et al. Human  
740 metabolic individuality in biomedical and pharmaceutical research. *Nature.* 2011;477:  
741 54–60.
- 742 33. Aulchenko YS, Struchalin MV, van Duijn CM. ProbABEL package for genome-wide  
743 association analysis of imputed data. *BMC Bioinformatics.* 2010;11: 134.

744

745 **TABLES**

746

747 **Table 1.** Study characteristics of the 694 females from the TwinsUK Cohort analyzed in this  
748 study.

749

Variable	Mean	SD	IQR	N
Age	58.51	8.90	12.02	694
Hours fasting	10.42	4.74	9.00	694
Time of day	11.75	2.06	4.00	694
Glucose	4.91	0.42	0.60	656
Insulin	44.20	23.76	31.00	622
Total Cholesterol	5.57	1.03	1.40	678
HDL	1.81	0.43	0.61	675
Triglycerides	1.01	0.39	0.54	647
Height	161.43	6.04	8.70	681
Weight	69.37	14.12	16.40	681
BMI	26.59	5.08	5.88	681

750

751

752

753

**Table 2 Number of genes associated with hours of fasting or time of day in adipose and skin tissue.** Each row lists the number of genes and exons whose expression is associated to the listed exposure in adipose tissue, skin tissue or both adipose and skin tissue respectively. The final row lists the number of genes that are associated to both hours of fasting and time of day in adipose tissue, skin tissue or both tissues.

	Adipose		Skin		Both tissues	
Exposure	Genes	Exons	Genes	Exons	Genes	Exons
Hours fasting	367	871	79	179	29	102
Time of day	99	211	54	127	17	67
Both exposures	71	150	19	73	15	56

764 **Table 3.** Magnitude of the effect size across tissues and exposures  
765

Exposure	Mean $ \beta $		S.D.		Cross - Tissue	Common Exons
	adipose	Skin	adipose	Skin	P	$ \beta_{\text{adipose}}  -  \beta_{\text{skin}} $
Hours fasting	0.048	0.046	0.011	0.009	1.2E-03	0.017
Time of day	0.107	0.106	0.017	0.025	7.0E-01	0.001
Cross-Exposure P	7.1E-131	5.3E-58	N.A.	N.A	N.A.	N.A.

766 \* Average magnitude of the difference between tissues of the regression coefficient for the  
767 subset of exons significant in both tissues  
768  
769



770 **Table 4.** Fasting or Time of day response genes with regulatory variants overlapping GWAS  
771 signals of complex traits.

Tissue	Exposure	GeneName	eQTL SNP	GWAS SNP	r2	Trait
Fat	Hours fasting	<i>ADAM19</i>	rs11134789	rs2277027	0.99	Lung function FEV1/FVC Lung function FEV1/FVC in ever smokers
Fat	Hours fasting	<i>CTPS1</i>	rs12037263	rs2802551	0.98	Height
Fat	Hours fasting	<i>DNM3</i>	rs17277932	rs17277932	1.00	Height
Fat	Hours fasting	<i>HEMK1</i>	rs399484	rs399484	1.00	Height
Fat	Hours fasting	<i>PHLDB1</i>	rs10790255	rs45565037	0.86	Rheumatoid arthritis
				rs10440635	1.00	Crohns disease
Fat	Hours fasting	<i>PTGER4</i>	rs10440635	rs7720838	1.00	inflammatory bowel disease Irritable bowel syndrome Self reported allergy
Fat	Hours fasting	<i>TRIM27</i>	rs115864834	rs429369	1.00	Height
Fat	Hours fasting	<i>TSEN15</i>	rs1952256	rs1046934	1.00	Height
Fat	Hours fasting & Time of day	<i>SLC9A3R2</i>	rs3211995	rs3211995	1.00	Height
Fat	Time of day	<i>TNFAIP8</i>	rs11064	rs1045241	0.99	Waist hip ratio in females Waist to hip ratio adjusted for body mass index
Skin	Hours fasting	<i>TNNT2</i>	rs12564445	rs12564445	1.00	Cardiac Troponin T levels

772

773 **Table 5.** Variants interacting with hours of fasting or time of day in fat and skin regulate  
774 expression of fasting or time of day associated genes.

Tissue	Exposure	Gene	SNP	Chr	P	FWER
Fat	Fasting	<i>ACTA2</i>	rs813782	10	9.8E-06	1.0E-02
Fat	Fasting	<i>CBFA2T3</i>	rs12921348	16	5.6E-05	3.4E-02
Fat	Fasting	<i>FRMD4A</i>	rs10906711	10	2.9E-05	4.3E-02
Fat	Fasting	<i>HK2</i>	rs11690080	2	3.7E-05	3.0E-02
Fat	Fasting	<i>KIAA0182</i>	rs56377278	16	4.9E-05	4.7E-02
Fat	Fasting	<i>KLHL25</i>	rs11854452	15	1.9E-05	1.8E-02
Fat	Fasting	<i>MBD5</i>	rs13016227	2	3.0E-05	1.0E-02
Fat	Fasting	<i>PLXND1</i>	rs2403343	3	1.0E-04	2.9E-02
Fat	Fasting	<i>PPFIBP1</i>	rs405394	12	6.0E-05	4.8E-02
Fat	Fasting	<i>RP11-399O19</i>	rs813782	10	7.6E-05	4.4E-02
Fat	Fasting	<i>STARD13</i>	rs9591423	13	8.4E-05	4.7E-02
Fat	Fasting	<i>THRA</i>	rs11651987	17	2.4E-05	1.3E-02
Fat	TOD	<i>FZD10</i>	rs11061011	12	4.0E-05	3.4E-02
Skin	TOD	<i>HR</i>	rs10110714	8	7.6E-05	4.8E-02

775  
776  
777

## **Figure Legends**

**Figure 1.** Overall distribution of (A) hours fasting, (B) hours fasting stratified by AM/PM, and (C) biopsy time of day in the 625 individuals participating in this study.

**Figure 2** Histogram of the P values for the association between expression levels of all expressed genes in adipose and skin tissue with time of the day (TOD) and hours fasting (HF): A) TOD in skin, B) TOD in adipose, C) HF in skin, HF in adipose.

**Figure 3. Tissue-specificity of gene expression response to hours of fasting and time of day.** Venn diagram of genes associated to hours fasting (top) and time of day (bottom) in adipose tissue (red) and skin (yellow) tissue. Genes associated to an exposure in both tissues are listed in the tables on the right. Tissue-shared genes are annotated to indicate genes that are members of the circadian clock or known to be regulated by circadian clock genes.

**Figure 4. Expression levels of most genes involved in the circadian clock are associated with fasting in adipose tissue.** Diagram shows the Ingenuity canonical pathway of the circadian clock annotated to highlight genes showing fasting-associated changes in fat gene expression in this study. Proteins (homo or heterodimer) with all coding genes associated to fasting are annotated in red, proteins with at least one gene associated with fasting are annotated with a pink border.

**Figure 5. SNP x Hours of Fasting interactions on gene expression.** The vertical axis represents expression of the the given gene and the horizontal axis hours of fasting. Each point represents one individual. Each plot is split into three panels with individuals plotted in the panel corresponding to their genotype at the given SNP. Solid line represents the linear

806 regression of expression on hours fasting, discontinuous line represents a smooth loess  
807 local regression

808  
809  
810  
811

812 **Additional File 1 : Alves\_AdditionalFile1.xlsx**

813 Additional File 1 is an excel worksheet containing 13 supplementary tables

814

815 **Supplementary Table S1.** Genes significantly associated with hours of fasting, time of day  
816 or seasonality in fat or skin

817

818 **Supplementary Table S2.** Replicability of literature studies and our study across taxa

819

820 **Supplementary Table S3.** Replicability of genes associated with hours fasting and time of  
821 day

822

823 **Supplementary Table S4.** Gene ontology enrichment of the hours fasting and solar time in  
824 fat and skin using TopGO redundancy elimination algorithms. FDR calculated using  
825 Benjamini Hochberg. Only GO Terms with < 500 annotated genes reported. Genes with  
826 association  $P < 1E-5$  considered significant.

827

828 **Supplementary Table S5.** Gene ontology enrichment of the hours fasting and solar time in  
829 fat and skin using ToppFun database and cumulative hypergeometric distribution test. FDR  
830 calculated using Benjamini Hochberg. Only GO Terms with < 500 annotated genes reported.  
831 Genes with association  $P < 1E-5$  considered significant

832

833 **Supplementary Table S6.** Ingenuity canonical pathways enrichment of the hours fasting  
834 and solar time in fat and skin using Ingenuity software. FDR calculated using Benjamini  
835 Hochberg on all pathways. Genes with association  $P < 1E-5$  considered significant.

836

837 **Supplementary Table S7.** Sensitivity analysis of GO term enrichment using Cut-off  $P < 2E-$   
838  $6$  equivalent to BY  $FDR < 5\%$ . Gene ontology enrichment of the hours fasting and solar time  
839 in fat and skin using TopGO redundancy elimination algorithms. FDR calculated using  
840 Benjamini Hochberg. Only GO Terms with < 500 annotated genes reported. Genes with  
841 association  $P < 1E-5$  considered significant.

842

843 **Supplementary Table S8.** Sensitivity analysis of GO term enrichment using Cut-off  $P < 5E-$   
844  $7$  equivalent to Bonferroni correction for 100, 000 independent tests. Gene ontology  
845 enrichment of the hours fasting and solar time in fat and skin using TopGO redundancy  
846 elimination algorithms. FDR calculated using Benjamini Hochberg. Only GO Terms with <  
847 500 annotated genes reported. Genes with association  $P < 1E-5$  considered significant.

848

849 **Supplementary Table S9.** Number of genes associated with hours fasting as a function of  
850 tissue, quantification, regression model. Significance cut-off  $P < 2E-6$ . FDR calculated using  
851 Benjamini-Hochberg approach.

**Supplementary Table S10.** Top pathway obtained with enrichment analysis of **Fat** and **Skin** gene expression as a function of quantification, regression model, and P-value corrections. Enrichment algorithm TopGO Fisher Elimination. Bonferroni-corrected significance  $\alpha_B=1\times 10^{-5}$  corresponding to approximately 5000 GO terms tested with size < 500 genes. LEGEND:




*\*Model gene p-value confounders:  $\text{logit}(P\text{-Value}) \sim \text{Intercept} + \text{GC percent} + \text{Gene length} + N \text{ effectively independent exons} + N \text{ of exons} + (1 \mid \text{Gene type}) + (1 \mid \text{gene status}) + (1 \mid \text{Database}) + (1 \mid \text{Strand})$ .*





*<sup>a</sup>The top 2 terms scored in ex equo the same P, the other term being “Negative regulation of transcription from RNA polymerase II promoter”. The third term is “Circadian regulation of gene expression” ( $P<7\times 10^{-7}$ ) with 8 significant out of 56 background genes and in the hierarchy a child of the most significantly enriched GO Term.*

**Supplementary Table S11.** Epidemiological identification of mediators of the response to fasting. Regression of the outcome on hours fasting adjusted for technical covariates, family structure and outcome-specific covariates identified glucose and total triglycerides as independent mediators of the metabolic response to fasting. LEGEND: *#The association of these outcomes with hours fasting was abrogated when conditioning on TG or Glucose, suggesting that TG and glucose might be confounders or mediators of the cardio-metabolic response to fasting.*


**Supplementary Table S12.** Genes with eQTL SNPs overlapping GWAS Signals associated with multiple phenotypes, enriched for cardio-metabolic and immune related phenotypes.

**Supplementary Table S13.** Genes with GxE eQTL SNPs with FWER<0.05.



**A****B****C****D**


## Hours Fasting



Adipose

Skin

## Time of Day



### Hours fasting tissue-overlapping genes

Circadian clock genes	Clock-regulated genes	Other genes
<i>ARNTL</i>	<i>DBP</i>	<i>APLN</i>
<i>BHLHE41</i>	<i>FKBP5</i>	<i>CAMKK1</i>
<i>C1orf51</i>	<i>ST3GAL5</i>	<i>CCND3</i>
<i>CRY2</i>	<i>TSC22D3</i>	<i>FAM46B</i>
<i>KLF9</i>	<i>UTS2</i>	<i>GPATCH11</i>
<i>NFIL3</i>	<i>ZBTB16</i>	<i>HHEX</i>
<i>NR1D1</i>		<i>MNT</i>
<i>NR1D2</i>		<i>RP3-67L1.4</i>
<i>PER1</i>		<i>SPHK2</i>
<i>PER2</i>		<i>THRA</i>
<i>PER3</i>		<i>TMCC3b</i>
<i>TEF</i>		

### Time of day tissue-overlapping genes

Circadian clock genes	Clock-regulated genes	Other genes
<i>ARNTL</i>	<i>FKBP5</i>	<i>FAM46B</i>
<i>C1orf51</i>	<i>HLF</i>	<i>SLC04A1</i>
<i>CRY2</i>	<i>TSC22D3</i>	
<i>KLF9</i>	<i>UTS2</i>	
<i>NR1D1</i>	<i>ZBTB16</i>	
<i>NR1D2</i>		
<i>PER1</i>		
<i>PER2</i>		
<i>PER3</i>		
<i>TEF</i>		



